




Contents List available at JACS Directory

Journal of Advanced Chemical Sciences

journal homepage: www.jacsdirectory.com/jacs

Stability Constants of Mixed Ligand Complexes of Transition Metal(II) ions with 1-{(E)-[(4-bromo-3-ethoxyphenyl)imino]methyl}naphthalen-2-ol and 4-bromo-2-{(E)-[(4-methoxy-2-methylphenyl)imino]methyl}phenol

A.K. Mapari*

Department of Chemistry, Ramnarain Ruia College, Matunga (E), Mumbai – 400 019, Maharashtra, India.

ARTICLE DETAILS

Article history:

Received 02 September 2016

Accepted 22 September 2016

Available online 13 November 2016

Keywords:

Binary Complexes

Ternary Complexes

Mixed Ligand

Stability Constant

ABSTRACT

Binary and ternary complexes of the type M-Y and M-X-Y [M = Co(II), Ni(II), Cu(II) and Zn(II); X = 4-bromo-2-{(E)-[(4-methoxy-2-methylphenyl)imino]methyl}phenol and Y = 1-{(E)-[(4-bromo-3-ethoxyphenyl)imino]methyl}naphthalen-2-ol] have been examined pH-metrically at 27 ± 0.5 °C and at constant ionic strength, $\mu = 0.1$ M (KCl) in 75:25 (v/v) 1,4-dioxane-water medium. The stability constants for Binary (M-Y) and ternary (M-X-Y) systems were calculated.

1. Introduction

Schiff bases play important roles in coordination chemistry as they easily form stable complexes with most transition metal ions [1, 2]. Many attempts have been made to evaluate different factors affecting the stability of the metal chelates along with their stability constants [3-5]. In the present study the stability constants of the mixed ligand complexes of Co(II), Ni(II), Cu(II) and Zn(II) with 4-bromo-2-{(E)-[(4-methoxy-2-methylphenyl)imino]methyl}phenol (X) as primary ligand (Fig. 1) and 1-{(E)-[(4-bromo-3-ethoxyphenyl)imino]methyl}naphthalen-2-ol (Y) as secondary ligand (Fig. 2) in 75:25 (v/v) 1,4-dioxane-water medium at 27 ± 0.5 °C have been reported by employing pH-metric titration technique [6-10]. Under identical conditions the stability constants of binary metal complexes of 1-{(E)-[(4-bromo-3-ethoxyphenyl)imino]methyl}naphthalen-2-ol (Y) have also been investigated.




Fig. 1 Primary ligand 4-bromo-2-{(E)-[(4-methoxy-2-methylphenyl)imino]methyl}phenol




Fig. 2 Secondary ligand 1-{(E)-[(4-bromo-3-ethoxyphenyl)imino]methyl}naphthalen-2-ol

2. Experimental Methods

The pH-meter model no. EQ-614 supplied by Equiptronics, a precision research pH-meter with wide range of glass electrode and calomel reference electrode was used for pH measurements. The pH-meter was standardized with potassium hydrogen phthalate and phosphate buffers before performing the titrations. The solutions of ligands were prepared in 1,4-dioxane. All the metal ion solutions were prepared in double distilled water and standardized by using conventional procedures [11]. A solution of KOH (0.2 M) was prepared in double distilled water and standardized with standard solution of succinic acid. The titrations were carried out in an inert atmosphere of nitrogen. All the measurements were carried out at temperature 27 ± 0.5 °C. The method of Bjerrum and Calvin as modified by Irving and Rossotti [6, 7] was used to determine $\bar{n}_A$ (average number of protons associated with secondary ligand); $\bar{n}$ (average number of secondary ligand molecules attached per metal ion); $\bar{n}_{mix}$ (average number of secondary ligand molecules attached per (M.X)⁺ ion); pL (free ligand exponent for binary (M-Y) system) and pL_{mix} (free ligand exponent for ternary (M-X-Y) system) values. All the solvents and chemicals used were of A R grade.

For the determination of proton-ligand stability constant of secondary ligand (Y) and metal-ligand stability constants of binary (M-Y) and ternary (M-X-Y) complexes, the following sets of solutions were prepared keeping the total volume $V_0 = 40$ mL. All titrations were carried out at the ionic strength of 0.1 M using KCl as an electrolyte in 75:25 (v/v) 1,4-dioxane-water medium against standard carbonate free KOH (0.2 M) solution.

- 4.0 mL HCl (0.16 M) + 3.36 mL KCl (1.0 M) + 2.64 mL distilled water + 30.0 mL 1,4-dioxane.
- 4.0 mL HCl (0.16 M) + 3.36 mL KCl (1.0 M) + 2.64 mL distilled water + 4.0 mL secondary ligand (0.04 M) + 26.0 mL 1,4-dioxane.
- 4.0 mL HCl (0.16 M) + 3.24 mL KCl (1.0 M) + requisite volume of metal(II) chloride solution to give 0.001 M metal(II) chloride concentration in the final solution + requisite amount of distilled water + 4.0 mL secondary ligand (0.04 M) + 26.0 mL 1,4-dioxane.
- 4.0 mL HCl (0.16 M) + 3.36 mL KCl (1.0 M) + 2.64 mL distilled water + 4.0 mL secondary ligand (0.01 M) + 26.0 mL 1,4-dioxane.
- 4.0 mL HCl (0.16 M) + 3.24 mL KCl (1.0 M) + requisite volume of metal(II) chloride solution to give 0.001 M metal(II) chloride concentration in the final solution + requisite amount of distilled water + 4.0 mL primary ligand (0.01 M) + 4.0 mL secondary ligand (0.01 M) + 22.0 mL 1,4-dioxane.

*Corresponding Author

Email Address: akmapari@gmail.com (A.K. Mapari)

The ratio of metal (M): secondary ligand (Y) was maintained at 1:4 in the binary system. In the ternary system the ratio of metal (M): primary ligand (X): secondary ligand (Y) was maintained at 1:1:1. The pH-meter readings were plotted against the volume of alkali (KOH) used for each titrations.

3. Results and Discussion

3.1 Proton-Ligand Stability Constants of Secondary Ligand (Y)

From the titration curves of solutions (i) and (ii), $\bar{n}$ A values at various pH were calculated. The proton ligand formation curve was obtained by plotting the values of $\bar{n}$ A vs. pH-meter readings. From the graph the values of $\log K_1^H$ and $\log K_2^H$ were evaluated by half integral method (A). The values of $\log K_1^H$ and $\log K_2^H$ were also evaluated using graphical method (B) by plotting the graph of $\log [\bar{n}A / (1-\bar{n}A)]$ against pH and $\log [(2-\bar{n}A) / (\bar{n}A-1)]$ against pH, respectively. The values obtained by method A and B are in agreement with each other, the average values of $\log K_1^H$ and $\log K_2^H$ has been found to be 8.65 and 2.11 respectively.

3.2 Metal-Ligand Stability Constants of the Binary (M-Y) Complexes

The metal ligand stability constants of binary complexes were evaluated assuming the polynuclear complexes and hydrolyzed products were not formed. The metal-ligand binary binding proposed structure is shown in Fig. 3. An examination of titration curves indicate that complex formation takes place in the solution on the following grounds: (I) The metal titration curve of solution (iii) shows displacement with respect ligand (Y) titration curve of solution (ii) along the volume axis. This indicates the affinity of the ligand to metal ions which release proton and produce volume difference. (II) The color change of ligand appeared in the presence of metal ion shows the formation of new species due to coordination. (III) The hydrolysis of the metal ion was suppressed due to the complex formation and precipitation did not appear during the titrations.




Fig. 3 The metal-ligand binary binding proposed structure

From the titration curves of solutions (ii) and (iii), $\bar{n}$ and pL values were calculated. The formation curves were obtained by plotting the values of $\bar{n}$ vs. pL. From the graph the values of $\log K_{MY_1}^M$ and $\log K_{MY_2}^{MY}$ were evaluated by half integral method (A) and in the similar way the values of $\log K_{MY_1}^M$ and $\log K_{MY_2}^{MY}$ were evaluated using graphical method (B) by plotting the graph of $\log [\bar{n} / (1-\bar{n})]$ against pL and $\log [(2-\bar{n}) / (\bar{n}-1)]$ against pL respectively. The values obtained by method A and B are in agreement with each other, the average values of $\log K_{MY_1}^M$ and $\log K_{MY_2}^{MY}$ along with metal-ligand stability constants the $\log \beta$ values of the binary complexes are given in Table 1.

Table 1 Metal-ligand stability constants of binary (M-Y), ternary (M-X-Y) systems and $\Delta \log K_T$ values in 75: 25 (v/v) 1,4-dioxane-water medium at 27 ± 0.5 °C and $\mu = 0.1$ M (KCl)

System	Metal-ligand stability constants / $\Delta \log K_T$	Metal(II) ions (M)			
		Co(II)	Ni(II)	Cu(II)	Zn(II)
Binary (M-Y)	$\log K_{MY_1}^M$	5.37	5.65	7.41	4.27
	$\log K_{MY_2}^{MY}$	3.73	3.63	5.65	3.32
	$\log \beta$	9.01	9.28	13.06	7.59
Ternary (M-X-Y)	$\log K_{MX}^{MX}$	5.18	5.26	7.23	4.03
(M-Y) and (M-X-Y)	$\Delta \log K_T$	-0.19	-0.39	-0.18	-0.24

The variations of $\bar{n}$ was found to be between 0.0-2.0 for the binary (M-Y) complexes of Co(II), Ni(II), Cu(II) and Zn(II) metal ions, which indicate that the composition of the complexes were 1:2 in solution. The $\log K_{MY_1}^M$ values for the binary complexes of the metal ions are in the following order: Cu(II) > Ni(II) > Co(II) > Zn(II), M = Co(II), Ni(II), Cu(II) and Zn(II).

3.3 Metal-Ligand Stability Constants of the Ternary (M-X-Y) Complexes

The metal-ligand stability constants of ternary complexes were evaluated assuming that the formation of polynuclear complexes and hydrolyzed products were not formed. The metal-ligand ternary binding proposed structure is shown in Fig. 4. An examination of the titration curves indicate that the ternary complexes formation has taken place in the solution on the following grounds: (I) the horizontal distance was measured between ternary titration curves of solution (v) and secondary ligand titration curve of solution (iv), the positive difference shows the earlier release of protons in the formation of ternary complexes. (II) The hydrolysis of the metal ion was suppressed and precipitation did not appear during the titrations.




Fig. 4 The metal-ligand ternary binding proposed structure

4. Conclusion

The value of $\log K_{MX}^{MX}$ are slightly lower than $\log K_{MY_1}^M$ and higher than $\log K_{MY_2}^{MY}$, which is due to the fact that the tendency of the secondary ligand (Y) to get bound with aquated metal ion $[M(aq)]^{2+}$ is more than to combine with the metal ion already bound with primary ligand (X). The relative stability ($\Delta \log K_T$) values of the ternary complexes with corresponding binary complexes for all the metal(II) ions in the present study is negative indicating that ternary 1:1:1 (M-X-Y) complexes are less stable than binary 1:1 (M-Y) complexes. In the ternary system studied, the order of stability constants of mixed ligand complexes with respect to the metal ions was found to be Cu(II) > Ni(II) > Co(II) > Zn(II); which is same as in the corresponding binary (M-Y) systems.

References

- [1] C. Spinu, M. Pleniceanu, C. Tigae, Biologically active transition metal chelates with a 2-Thiophenecarboxaldehyde-derived Schiff base: synthesis, characterization, and antibacterial properties, J. Turk. Chem. 32 (2008) 487-493.
- [2] B. Clarke, N. Clarke, D. Cunningham, T. Higgins, P. McArdle, et al, Transition-metal schiff-base complexes as ligands in tin chemistry part 7 reactions of organotin(IV) lewis acids with $[M(L)]_2$ [M-Ni, Cu and Zn; H₂L-N,N-bis(3-methoxysalicylidene)benzene-1,3-diamine and its -1,4-diamine analog], J. Organometallic Chem. 559 (1998) 55-64.
- [3] C.N. Mukherjee, T. Ghosh, Mixed ligand complex formation of Cobalt(II), Nickel(II) and Zinc(II) with N-Benzenesulphonyl-L-cysteine as primary ligand and Bipyridine, Ethylenediamine and Glycinate as secondary ligands, J. Ind. Chem. Soc. 74 (1977) 8-11.
- [4] K. Kiramany, Y. Prashanthi, N.J.P. Subhashini, Shivraj, Synthesis, characterization and biological activity of metal complexes of 3-amino-5-methyl isoxazole Schiff bases, J. Chem. Pharm. Res. 2(1) (2010) 375-384.
- [5] B.C. Khade, P.M. Deore, B.R. Arbad, Mixed-ligand complex formation of copper (II) with some amino acids and Drug dapsone, Int. J. Chem. Tech. Res. 2(2) (2010) 1036-1040.
- [6] H.M. Irving, H.S. Rossotti, Methods for computing successive stability constants from experimental formation curves, J. Chem. Soc. (1953) 3397-3405.
- [7] H.M. Irving, H.S. Rossotti, The Calculations of formations curves of metal complexes from pH titration curves in mixed solvents, J. Chem. Soc. (1954) 2904-2910.

- [8] S. Valecha, K. Mangaonkar, Mixed ligand chelates of transition metal ions with [1-Phenyl-3-methyl-5-oxodihydropyrazole-4-yl] phenylimino]-2', 3'-dimethylaniline as a primary ligand and 2-Hydroxy-1-naphthaldehyde as a secondary ligand, *Asian J. Chem.* 20(2) (2008) 967-972.
- [9] V. Rukmani, K. Prachi K.V. Mangaonkar, Mixed ligand chelates of transition metal ions with 5-Chlorosalicylidene-2-chlorobenzylamine as a primary ligand and 5-Chlorosalicylidene-4-aminobenzenesulphonamide as a secondary ligand, *Asian J. Chem.* 20(2) (2008) 949-953.
- [10] Ghasemi, Lahanbakhsh, Shokrollahi, Potentiometric study of binary and mixed complexes of imidazole, histamine, histidine and diacetyl monooxime with some transition metal ions in aqueous solution, Iran. J. Chem. Chem. Engg. 20(1) (2001) 22-28.
- [11] A.I. Vogel, Text book of quantitative practical inorganic chemistry, ELBS, England, 1984.
- [12] A.K. Patel, J.D. Joshi, Potentiometric studies on mixed ligand complexes in aqueous solution: M(II) + neutral or charged primary ligand + amino acid systems, *J. Ind. Chem. Soc.* 74 (1997) 222-224.
- [13] S. Satyanarayana, K.V. Reddy, stability constants of the binary complexes of Co(II), Ni(II), Cu(II) and Zn(II) with thiamine and ternary complexes with thiamine as primary ligand and thymine or uracil as secondary ligand, *Ind. J. Chem.* 28(A) (1989) 169-171.